

Package ‘MetaGxBreast’

June 12, 2025

Type Package

Title Transcriptomic Breast Cancer Datasets

Version 1.28.0

Date 2020-04-23

Description A collection of Breast Cancer Transcriptomic Datasets that are part of the MetaGxData package compendium.

License Apache License (>= 2)

Depends R (>= 3.6.0), Biobase, AnnotationHub, ExperimentHub

Imports stats, lattice, impute, SummarizedExperiment

Suggests testthat, xtable, tinytex

NeedsCompilation no

biocViews ExpressionData, ExperimentHub, CancerData,
Homo_sapiens_Data, ArrayExpress, GEO, NCI, MicroarrayData,
ExperimentData

LazyData yes

RoxygenNote 7.1.1

git_url <https://git.bioconductor.org/packages/MetaGxBreast>

git_branch RELEASE_3_21

git_last_commit 9e95d03

git_last_commit_date 2025-04-15

Repository Bioconductor 3.21

Date/Publication 2025-06-12

Author Michael Zon [aut],
Deena M.A. Gendoo [aut],
Christopher Eeles [ctb],
Benjamin Haibe-Kains [aut, cre]

Maintainer Benjamin Haibe-Kains <benjamin.haibe.kains@utoronto.ca>

Contents

CAL	3
DFHCC	5
DFHCC2	8
DFHCC3	10
DUKE	11
DUKE2	13
duplicates	15
EMC2	16
EORTC10994	18
EXPO	20
FNCLCC	22
GSE25066	23
GSE32646	26
GSE48091	28
GSE58644	29
HLP	32
IRB	34
KOO	36
loadBreastDatasets	38
loadBreastEsets	39
LUND	40
LUND2	42
MAINZ	44
MAQC2	46
MCCC	48
MDA4	49
METABRIC	51
MSK	54
MUG	56
NCCS	57
NCI	59
NKI	61
PNC	64
STK	66
STNO2	68
TCGA	71
TRANSBIG	73
UCSF	75
UNC4	78
UNT	81
UPP	83
VDX	85

 CAL

 CAL

Description

ExpressionSet for the CAL Dataset

Format

```

experimentData(eset):
Experiment data
  Experimenter name:
  Laboratory:
  Contact information: http://www.ncbi.nlm.nih.gov/pubmed/17157792
  Title:
  URL: http://www.ebi.ac.uk/arrayexpress/experiments/E-TABM-158/
  PMIDs: 17157792
  No abstract available.
  notes:
    summary:
      Recurrent copy number abnormalities differ between tumor subtypes as defin
ed by gene expression patterns. Accuracy of stratification by outcome can
be improved by combining expression and copy number.
    mapping.method:
      maxRowVariance
    mapping.group:
      EntrezGene.ID
    preprocessing:
      As published by original author.

featureData(eset):
An object of class 'AnnotatedDataFrame'
  featureNames: 1007_s_at 1053_at ... AFFX-HUMISGF3A/M97935_MB_at
  (21169 total)
  varLabels: probeset gene EntrezGene.ID best_probe
  varMetadata: labelDescription

```

Details

```

assayData: 21169 features, 118 samples
Platform type:
Overall survival time-to-event summary (in years):
Call: survfit(formula = Surv(time, cens) ~ -1)

1 observation deleted due to missingness
  n events median 0.95LCL 0.95UCL

```

117.00 77.00 8.96 8.33 9.71

Available sample meta-data:

sample_name:

Length	Class	Mode
118	character	character

sample_type:

tumor
118

er:

negative	positive
43	75

pgr:

negative	positive	NA's
51	66	1

tumor_size:

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.	NA's
0.300	1.675	2.300	2.729	3.500	7.500	2

N:

0	1
51	67

age_at_initial_pathologic_diagnosis:

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.	NA's
31.00	44.00	51.00	55.06	66.00	88.00	1

grade:

1	2	3	NA's
10	42	61	5

dmfs_days:

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.	NA's
0	767	2059	2094	3336	5183	1

dmfs_status:

norecurrence	recurrence	NA's
91	26	1

days_to_tumor_recurrence:

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.	NA's
------	---------	--------	------	---------	------	------

```

0      767      2059      2094      3336      5183      1

recurrence_status:
norecurrence  recurrence      NA's
           81           36           1

days_to_death:
  Min. 1st Qu.  Median    Mean 3rd Qu.    Max.    NA's
    47   1117   2234   2347   3504   5183      1

vital_status:
deceased  living
      77      41

treatment:
chemo.plus.hormono      chemotherapy      hormonotherapy      untreated
           25           36           40           14
      NA's
           3

batch:
CAL
118

uncurated_author_metadata:
  Length    Class      Mode
    118 character character

```

Source

<http://www.ebi.ac.uk/arrayexpress/experiments/E-TABM-158/>

DFHCC

DFHCC

Description

ExpressionSet for the DFHCC Dataset

Format

```

experimentData(eset):
Experiment data
  Experimenter name:
  Laboratory:
  Contact information: http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2826790/

```

```

Title:
URL: http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE19615
PMIDs: 20098429
No abstract available.
notes:
  summary:
    A small number of over-expressed and over-amplified genes were significant
ly associated with early recurrence despite adjuvant therapy. This was ve
rified in independent cohorts.
  mapping.method:
    maxRowVariance
  mapping.group:
    EntrezGene.ID
  preprocessing:
    As published by original author.

featureData(eset):
An object of class 'AnnotatedDataFrame'
  featureNames: 1007_s_at 1053_at ... AFFX-HUMISGF3A/M97935_MB_at
    (42447 total)
  varLabels: probeset gene EntrezGene.ID best_probe
  varMetadata: labelDescription

```

Details

```

assayData: 42447 features, 115 samples
Platform type:
-----
Available sample meta-data:
-----

sample_name:
  Length      Class      Mode
    115 character character

alt_sample_name:
  Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
   6.0   155.0   230.0   293.3   398.5   828.0

sample_type:
tumor
  115

er:
negative positive
   45         70

pgr:

```

```

negative positive
      51      64

```

```

her2:
negative positive
      79      36

```

```

tumor_size:
  Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
  0.800  1.350  2.100  2.312  2.850  6.500

```

```

N:
  0  1
62 53

```

```

age_at_initial_pathologic_diagnosis:
  Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
  32.00  45.00  53.00  53.89  60.00  85.00

```

```

grade:
  1  2  3
23 28 64

```

```

dmfs_days:
  Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
    30   1500   1920   1799   2325   2640

```

```

dmfs_status:
norecurrence  recurrence
          101           14

```

```

treatment:
chemo.plus.hormono      chemotherapy      hormonotherapy      untreated
              42              38              22              7
              NA's
              6

```

```

batch:
DFHCC
  115

```

```

uncurated_author_metadata:
  Length    Class      Mode
    115 character character

```

Source

<http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE19615>

DFHCC2

DFHCC2

Description

Test the efficacy of treating TNBC with neoadjuvant cisplatin; explore biomarkers to identify predictors of response

Format

```
experimentData(eset):
```

```
Experiment data
```

```
  Experimenter name:
```

```
  Laboratory:
```

```
  Contact information: http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2834466/
```

```
  Title:
```

```
  URL: http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE18864
```

```
  PMIDs: 20100965
```

```
Abstract: A 16 word abstract is available. Use 'abstract' method.
```

```
notes:
```

```
  summary:
```

```
    A subset of the patients experienced a response induced by cisplatin and b
    iomarkers were identified that could predict response to cisplatin.
```

```
  mapping.method:
```

```
    maxRowVariance
```

```
  mapping.group:
```

```
    EntrezGene.ID
```

```
  preprocessing:
```

```
    As published by original author.
```

```
featureData(eset):
```

```
An object of class 'AnnotatedDataFrame'
```

```
  featureNames: 1007_s_at 1053_at ... AFFX-HUMISGF3A/M97935_MB_at
  (42447 total)
```

```
  varLabels: probeset gene EntrezGene.ID best_probe
```

```
  varMetadata: labelDescription
```

Details

```
assayData: 42447 features, 84 samples
```

```
Platform type:
```

```
-----
```

```
Available sample meta-data:
```

```
-----
```

```

sample_name:
  Length      Class      Mode
    84 character character

unique_patient_ID:
  Length      Class      Mode
    84 character character

sample_type:
tumor
  84

er:
negative positive
  53      31

pgr:
negative positive
  53      31

her2:
negative positive
  66      18

age_at_initial_pathologic_diagnosis:
  Min. 1st Qu.  Median    Mean 3rd Qu.  Max.
  29.00  45.00   53.00   52.89  59.00   85.00

grade:
  1  2  3
10 16 58

treatment:
chemotherapy
  84

batch:
DFHCC2_CISPLATIN DFHCC2_REFERENCE
                24                60

uncurated_author_metadata:
  Length      Class      Mode
    84 character character

duplicates:
  Length      Class      Mode
    84 character character

```

Source

<http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE18864>

DFHCC3

*DFHCC3***Description**

ExpressionSet for the DFHCC3 Dataset

Format

```

experimentData(eset):
Experiment data
  Experimenter name:
  Laboratory:
  Contact information: http://www.ncbi.nlm.nih.gov/pubmed/16473279
  Title:
  URL: http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE3744
  PMIDs: 16473279
  No abstract available.
  notes:
    summary:
      Basal like cancerse often lack an inactivated X chromosome.e_Other markers
      found were duplication of the active X chromosome ande_nonheterochromatin
      ized X chromosomal DNA. A small subset of X chromosomal genes were overexp
      ressed. These abnormalities are thought to led to the pathogenesis of basa
      l like cancers.
    mapping.method:
      maxRowVariance
    mapping.group:
      EntrezGene.ID
    preprocessing:
      As published by original author.

featureData(eset):
An object of class 'AnnotatedDataFrame'
  featureNames: 1007_s_at 1053_at ... AFFX-HUMISGF3A/M97935_MB_at
    (42447 total)
  varLabels: probeset gene EntrezGene.ID best_probe
  varMetadata: labelDescription

```

Details

assayData: 42447 features, 40 samples
 Platform type:

```

-----
Available sample meta-data:
-----

sample_name:
  Length      Class      Mode
    40 character character

alt_sample_name:
  Length      Class      Mode
    40 character character

sample_type:
tumor
    40

batch:
DFHCC3
    40

uncurated_author_metadata:
  Length      Class      Mode
    40 character character

```

Source

<http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE3744>

DUKE

DUKE

Description

ExpressionSet for the DUKE Dataset

Format

```

experimentData(eset):
Experiment data
  Experimenter name:
  Laboratory:
  Contact information: http://www.ncbi.nlm.nih.gov/pubmed/16273092
  Title:
  URL: http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=gse3143
  PMIDs: 16273092
  No abstract available.

```

```

notes:
  summary:
    It was shown that the activatione_status of several oncogenic pathways can
    bee_identified by gene expression signatures. These gene signatures ident
    ify deregulation of pathways, associations with clinically relevant outcom
    es, and characteristics of specific cancers and tumor subtypes.
  mapping.method:
    maxRowVariance
  mapping.group:
    EntrezGene.ID
  preprocessing:
    As published by original author.

featureData(eset):
An object of class 'AnnotatedDataFrame'
  featureNames: 1000_at 1001_at ... AFFX-MurIL4_at (12085 total)
  varLabels: probeset gene EntrezGene.ID best_probe
  varMetadata: labelDescription

```

Details

```

assayData: 12085 features, 171 samples
Platform type:
Overall survival time-to-event summary (in years):
Call: survfit(formula = Surv(time, cens) ~ -1)

```

```

  1 observation deleted due to missingness
      n  events  median 0.95LCL 0.95UCL
170.00  43.00   9.01   6.22    NA

```

```

-----
Available sample meta-data:
-----

```

```

sample_name:
  Length      Class      Mode
    171 character character

```

```

alt_sample_name:
  Length      Class      Mode
    171 character character

```

```

sample_type:
tumor
  171

```

```

er:
negative positive

```

```

      57      114

pgr:
negative positive  NA's
      23      65      83

tumor_size:
  Min. 1st Qu.  Median    Mean 3rd Qu.    Max.   NA's
  0.20   1.80   2.30   2.74   3.50   8.50    83

N:
  0      1 NA's
 53   36   82

days_to_death:
  Min. 1st Qu.  Median    Mean 3rd Qu.    Max.   NA's
 171.0  417.0  957.5 1235.0 1852.0 4069.0     1

vital_status:
deceased  living    NA's
      43     127      1

batch:
DUKE
171

uncurated_author_metadata:
  Length      Class      Mode
    171 character character

duplicates:
DUKE.DUKE_T00.622 DUKE.DUKE_T01.052 DUKE.DUKE_T01.522 DUKE.DUKE_T01.534
              1              1              1              1
              NA's
              167
```

Source

<http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=gse3143>

DUKE2	<i>DUKE2</i>
-------	--------------

Description

Predicting response with gene signature

Format

```

experimentData(eset):
Experiment data
  Experimenter name:
  Laboratory:
  Contact information: http://www.ncbi.nlm.nih.gov/pubmed/18024211
  Title:
  URL: http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=gse6861
  PMIDs: 18024211

  Abstract: A 5 word abstract is available. Use 'abstract' method.
  notes:
    summary:
      Retraction in Lancet Feb 2011 (21277543); Regimen specific signatures were
      able to predict pathological complete response. Selecting patients with t
      hese gene signataures could increase the proportion of patients with pCR t
      han by basing clinical decisions on clinical factors.
    mapping.method:
      maxRowVariance
    mapping.group:
      EntrezGene.ID
    preprocessing:
      As published by original author.

featureData(eset):
An object of class 'AnnotatedDataFrame'
  featureNames: 1053_3p_at 117_3p_at ... X79510cds_3p_s_at (45490
    total)
  varLabels: probeset gene EntrezGene.ID best_probe
  varMetadata: labelDescription

```

Details

```

assayData: 45490 features, 160 samples
Platform type:
-----
Available sample meta-data:
-----

sample_name:
  Length      Class      Mode
    160 character character

alt_sample_name:
  Length      Class      Mode
    160 character character

```

```
sample_type:
tumor
160

er:
negative positive
123      37

pgr:
negative positive NA's
133      25      2

N:
  0      1 NA's
58  95      7

age_at_initial_pathologic_diagnosis:
  Min. 1st Qu.  Median    Mean 3rd Qu.    Max.    NA's
26.00  43.00  49.00  49.41  56.00  70.00     35

grade:
  1      2      3 NA's
  2  37  70  51

treatment:
chemotherapy
160

batch:
DUKE2
160

uncurated_author_metadata:
  Length      Class      Mode
    160 character character
```

Source

<http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=gse6861>

duplicates	<i>a list containing the names of patients that are believed to be duplicates across datasets</i>
------------	---

Description

The object is a list where each element is a patient ID that is believed to be a duplicate of a patient in another dataset. Patients are designated as duplicated if they have Spearman correlations greater than or equal to 0.98 with other patient expression profiles

Format

A list with 107 elements, each of which is a patient ID.

EMC2

EMC2

Description

ExpressionSet for the EMC2 Dataset

Format

```
experimentData(eset):
Experiment data
  Experimenter name:
  Laboratory:
  Contact information: http://www.ncbi.nlm.nih.gov/pubmed/19421193
  Title:
  URL: http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=gse12276
  PMIDs: 19421193
  No abstract available.
  notes:
    summary:
      Genes were identified that may increase the ability of breast cancer cells
      to infiltrate the blood-brain barrier.
    mapping.method:
      maxRowVariance
    mapping.group:
      EntrezGene.ID
    preprocessing:
      As published by original author.

featureData(eset):
An object of class 'AnnotatedDataFrame'
  featureNames: 1007_s_at 1053_at ... AFFX-HUMISGF3A/M97935_MB_at
  (42447 total)
  varLabels: probeset gene EntrezGene.ID best_probe
  varMetadata: labelDescription
```

Details

assayData: 42447 features, 204 samples

Platform type:

Available sample meta-data:

sample_name:

Length	Class	Mode
204	character	character

alt_sample_name:

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
1.00	51.75	102.50	102.50	153.20	204.00

sample_type:

tumor
204

N:

0 NA's
48 156

dmfs_days:

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
0	335	640	799	1098	3507

dmfs_status:

norecurrence	recurrence
19	185

treatment:

chemotherapy	untreated
156	48

batch:

EMC2
204

uncurated_author_metadata:

Length	Class	Mode
204	character	character

Source

<http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=gse12276>

EORTC10994

*EORTC10994***Description**

ExpressionSet for the EORTC10994 Dataset

Format

```

experimentData(eset):
Experiment data
  Experimenter name:
  Laboratory:
  Contact information: http://www.ncbi.nlm.nih.gov/pubmed/?term=15897907
  Title:
  URL: http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE1561
  PMIDs: 15897907
  No abstract available.
  notes:
    summary:
      The tumors with an apocrine gene expression profile had strong histologica
l apocrine features. These tumors were androgen receptor positive and were
all ER negative, creating further classifications of tumor cells based on
steroid receptor activity- luminal which are ER and AR positive, basal th
at are ER and AR negative, and molecular apocrine that are ER negative and
AR positive.
    mapping.method:
      maxRowVariance
    mapping.group:
      EntrezGene.ID
    preprocessing:
      As published by original author.

featureData(eset):
An object of class 'AnnotatedDataFrame'
  featureNames: 1007_s_at 1053_at ... AFFX-HUMISGF3A/M97935_MB_at
  (20967 total)
  varLabels: probeset gene EntrezGene.ID best_probe
  varMetadata: labelDescription

```

Details

```

assayData: 20967 features, 49 samples
Platform type:
-----
Available sample meta-data:

```

```

-----

sample_name:
  Length      Class      Mode
    49 character character

alt_sample_name:
  Length      Class      Mode
    49 character character

sample_type:
tumor
  49

er:
negative positive
  22      27

pgr:
negative positive  NA's
  29      18      2

tumor_size:
  1  2  3  4
  4 23 14  8

N:
  0  1
19 30

grade:
  1  2  3 NA's
  4 22 20  3

batch:
EORTC10994
  49

uncurated_author_metadata:
  Length      Class      Mode
    49 character character

```

Source

<http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE1561>

EXPO

*EXPO***Description**

ExpressionSet for the EXPO Dataset

Format

```

experimentData(eset):
Experiment data
  Experimenter name:
  Laboratory:
  Contact information:
  Title:
  URL: http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE2109
  PMIDs:
  No abstract available.
  notes:
    summary:
      N/A
    mapping.method:
      maxRowVariance
    mapping.group:
      EntrezGene.ID
    preprocessing:
      As published by original author.

featureData(eset):
An object of class 'AnnotatedDataFrame'
  featureNames: 1007_s_at 1053_at ... AFFX-HUMISGF3A/M97935_MB_at
    (42447 total)
  varLabels: probeset gene EntrezGene.ID best_probe
  varMetadata: labelDescription

```

Details

```

assayData: 42447 features, 353 samples
Platform type:
-----
Available sample meta-data:
-----

sample_name:
  Length      Class      Mode
    353 character character

```

```

alt_sample_name:
  Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
  1005   21640  101100  134700  215900  486200

sample_type:
tumor
  353

er:
negative positive  NA's
   85      161     107

pgr:
negative positive  NA's
  114      129     110

her2:
negative positive  NA's
  166      61     126

age_at_initial_pathologic_diagnosis:
  Min. 1st Qu.  Median    Mean 3rd Qu.    Max.  NA's
  25.00  45.00  55.00  59.44  67.50  95.00    1

grade:
  1    2    3 NA's
  32 114 151  56

batch:
EXPO
  353

uncurated_author_metadata:
  Length    Class      Mode
  353 character character

duplicates:
EXPO.EXPO_GSM53027 EXPO.EXPO_GSM53059  NA's
                   1                   1    351

```

Source

<http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE2109>

FNCLCC

*FNCLCC***Description**

ExpressionSet for the FNCLCC Dataset

Format

```

experimentData(eset):
Experiment data
  Experimenter name:
  Laboratory:
  Contact information: http://www.ncbi.nlm.nih.gov/pubmed/?term=17659439
  Title:
  URL: http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE7017
  PMIDs: 17659439
  No abstract available.
  notes:
    summary:
      A potentially more powerful clinicogenomic model was created by combining
      a subset of relevant genes from an already published gene expression signa
      ture and a commonly used clinical prognostic model (NPI). The genes in thi
      s model are known to have a role in breast cancer, carcinogenesis, or chem
      otherapy resistance.
    mapping.method:
      maxRowVariance
    mapping.group:
      EntrezGene.ID
    preprocessing:
      As published by original author.

featureData(eset):
An object of class 'AnnotatedDataFrame'
  featureNames: UMGC_000005 UMGC_000007 ... UMGC_09018 (6064 total)
  varLabels: probeset gene EntrezGene.ID best_probe
  varMetadata: labelDescription

```

Details

```

assayData: 6064 features, 150 samples
Platform type:
-----
Available sample meta-data:
-----

```

```
sample_name:
  Length      Class      Mode
    150 character character

alt_sample_name:
  Length      Class      Mode
    150 character character

sample_type:
tumor
    150

N:
  1
    150

treatment:
chemotherapy
    150

batch:
FNCLCC
    150

uncurated_author_metadata:
  Length      Class      Mode
    150 character character
```

Source

<http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE7017>

GSE25066	<i>GSE25066</i>
----------	-----------------

Description

ExpressionSet for the GSE25066 Dataset

Format

```
experimentData(eset):
Experiment data
  Experimenter name:
  Laboratory:
  Contact information:
```

```

Title:
URL: http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE25066
PMIDs: 21558518
No abstract available.
notes:
  summary:

  mapping.method:
    maxRowVariance
  mapping.group:
    EntrezGene.ID
  preprocessing:
    As published by original author.

featureData(eset):
An object of class 'AnnotatedDataFrame'
  featureNames: 1007_s_at 1053_at ... AFFX-HUMISGF3A/M97935_MB_at
    (20967 total)
  varLabels: probeset gene EntrezGene.ID best_probe
  varMetadata: labelDescription

```

Details

assayData: 20967 features, 508 samples

Platform type:

 Available sample meta-data:

sample_name:

Length	Class	Mode
508	character	character

alt_sample_name:

Length	Class	Mode
508	character	character

sample_type:

tumor
508

er:

negative	positive	NA's
205	297	6

pgr:

negative	positive	NA's
258	243	7

```

her2:
negative positive  NA's
  485         6    17

T:
T0  T1  T2  T3  T4
  3  30 255 145  75

N:
  0  1
157 351

age_at_initial_pathologic_diagnosis:
  Min. 1st Qu.  Median    Mean 3rd Qu.  Max.
  24.0   42.0   49.0   49.8   58.0   75.0

grade:
  1  2  3  4 NA's
 32 180 259 15  22

dmfs_days:
  Min. 1st Qu.  Median    Mean 3rd Qu.  Max.
  0.0   636.5   999.9 1088.0 1500.0 2717.0

dmfs_status:
norecurrence  recurrence
      397         111

batch:
GSE25066
  508

uncurated_author_metadata:
  Length    Class      Mode
  508 character character

chemosensitivity_prediction:
Rx Insensitive  Rx Sensitive
      339         169

GGI_prediction:
High  Low
 336  172

PAM50_prediction:
Basal  Her2  LumA  LumB  Normal
  189    37   160    78    44

```

```

dlda30_prediction:
pCR  RD
196 312

RCB_prediction:
      RCB-0/I RCB-II/III
      230      278

```

Source

<http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE25066>

GSE32646

GSE32646

Description

ExpressionSet for the GSE32646 Dataset

Format

```

experimentData(eset):
Experiment data
  Experimenter name:
  Laboratory:
  Contact information:
  Title:
  URL: http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE32646
  PMIDs: 22320227
  No abstract available.
  notes:
    summary:

  mapping.method:
    maxRowVariance
  mapping.group:
    EntrezGene.ID
  preprocessing:
    As published by original author.

featureData(eset):
An object of class 'AnnotatedDataFrame'
  featureNames: 1007_s_at 1053_at ... 91952_at (42437 total)
  varLabels: probeset gene EntrezGene.ID best_probe
  varMetadata: labelDescription

```

Details

assayData: 42437 features, 115 samples

Platform type:

Available sample meta-data:

sample_name:

Length	Class	Mode
115	character	character

sample_type:

tumor

115

er:

negative positive

44 71

pgr:

negative positive

70 45

her2:

negative positive

81 34

T:

1 2 3 4

5 87 18 5

N:

0 1

32 83

age_at_initial_pathologic_diagnosis:

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
27.00	45.00	51.00	51.49	59.00	73.00

grade:

1 2 3

16 78 21

batch:

GSE32646

115

uncurated_author_metadata:

Length	Class	Mode
115	character	character

Source

<http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE32646>

GSE48091

GSE48091

Description

ExpressionSet for the GSE48091 Dataset

Format

```
experimentData(eset):
Experiment data
  Experimenter name:
  Laboratory:
  Contact information:
  Title:
  URL: http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE48091
  PMIDs: 26077471
  No abstract available.
  notes:
    summary:

    mapping.method:
      maxRowVariance
    mapping.group:
      EntrezGene.ID
    preprocessing:
      As published by original author.

featureData(eset):
An object of class 'AnnotatedDataFrame'
  featureNames: 100121619_TGI_at 100121620_TGI_at ... 100314044_TGI_at
    (23246 total)
  varLabels: probeset gene EntrezGene.ID best_probe
  varMetadata: labelDescription
```

Details

assayData: 23246 features, 623 samples

Platform type:

Available sample meta-data:

sample_name:

Length	Class	Mode
623	character	character

sample_type:

tumor

623

batch:

GSE48091

623

uncurated_author_metadata:

Length	Class	Mode
623	character	character

Source

<http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE48091>

GSE58644

GSE58644

Description

ExpressionSet for the GSE58644 Dataset

Format

experimentData(eset):

Experiment data

 Experimenter name:

 Laboratory:

 Contact information:

 Title:

 URL: <http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE58644>

 PMIDs: 25284793

 No abstract available.

 notes:

```

summary:

mapping.method:
  maxRowVariance
mapping.group:
  EntrezGene.ID
preprocessing:
  As published by original author.

featureData(eset):
An object of class 'AnnotatedDataFrame'
  featureNames: 7896756 7896759 ... 8180179 (21462 total)
  varLabels: probeset gene EntrezGene.ID best_probe
  varMetadata: labelDescription

```

Details

```

assayData: 21462 features, 321 samples
Platform type:

```

```

-----
Available sample meta-data:
-----

```

```

sample_name:
  Length      Class      Mode
    321 character character

```

```

alt_sample_name:
  Length      Class      Mode
    321 character character

```

```

sample_type:
tumor
  321

```

```

er:
negative positive  NA's
    70      250      1

```

```

her2:
negative positive  NA's
    256      58      7

```

```

tumor_size:
  Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
  0.600  1.600   2.100   2.354  2.600  15.000

```

```

T:

```

```

      1      2      3      4 NA's
43    59    13     1    205

N:
  0      1 NA's
138 151    32

age_at_initial_pathologic_diagnosis:
  Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
 29.00  49.00  58.00  58.82  68.00   93.00

grade:
  1      2      3 NA's
26 135 159     1

dmfs_status:
norecurrence  recurrence
          295           26

dmfs_days:
  Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
    0    9496  17900  21620  33600  52590

treatment:
chemo.plus.hormono      chemotherapy      hormonotherapy      untreated
              91              29              66              10
              NA's
              125

chemo:
  0      1 NA's
105 123  93

tamoxifen:
  0      1 NA's
39 157 125

herceptin:
  0      1 NA's
190  12 119

batch:
GSE58644
      321

uncurated_author_metadata:
  Length      Class      Mode
    321 character character
```

```

duplicates:
  Length      Class      Mode
    321 character character

```

Source

<http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE58644>

HLP

HLP

Description

ExpressionSet for the HLP Dataset

Format

```

experimentData(eset):
Experiment data
  Experimenter name:
  Laboratory:
  Contact information: http://www.ncbi.nlm.nih.gov/pubmed/?term=19688261
  Title:
  URL: http://www.ebi.ac.uk/arrayexpress/experiments/E-TABM-543/
  PMIDs: 19688261
  No abstract available.
notes:
  summary:
    The results show evidence of different patterns of genetic aberrations in
    distinct molecular subtypes of breast cancer. Patterns of copy number aber-
    rations may drive biological phenomena characteristic to each subtype.
  mapping.method:
    maxRowVariance
  mapping.group:
    EntrezGene.ID
  preprocessing:
    As published by original author.

featureData(eset):
An object of class 'AnnotatedDataFrame'
  featureNames: 9g8cQB1TZtuiix.ulU fJUdX0IAn_P9VLTgJU ...
    xopB7pPn18FJ067uDs (26536 total)
  varLabels: probeset gene EntrezGene.ID best_probe
  varMetadata: labelDescription

```

Details

assayData: 26536 features, 53 samples

Platform type:

Available sample meta-data:

sample_name:

Length	Class	Mode
53	character	character

alt_sample_name:

Length	Class	Mode
53	character	character

sample_type:

tumor

53

er:

negative positive

28	25
----	----

pgr:

negative positive

33	20
----	----

her2:

negative positive

40	13
----	----

tumor_size:

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.	NA's
1.200	1.800	2.450	2.648	3.000	8.000	5

N:

0	1	NA's
27	25	1

age_at_initial_pathologic_diagnosis:

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.	NA's
30.00	47.50	53.50	54.96	64.25	81.00	5

grade:

3

53

batch:

HLP
53

uncurated_author_metadata:
Length Class Mode
53 character character

Source

<http://www.ebi.ac.uk/arrayexpress/experiments/E-TABM-543/>

IRB	<i>IRB</i>
-----	------------

Description

ExpressionSet for the IRB Dataset

Format

```
experimentData(eset):
Experiment data
  Experimenter name:
  Laboratory:
  Contact information: http://www.ncbi.nlm.nih.gov/pubmed/18297396
  Title:
  URL: http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE5460
  PMIDs: 18297396
  No abstract available.
  notes:
    summary:

    mapping.method:
      maxRowVariance
    mapping.group:
      EntrezGene.ID
    preprocessing:
      As published by original author.

featureData(eset):
An object of class 'AnnotatedDataFrame'
  featureNames: 1007_s_at 1053_at ... AFFX-HUMISGF3A/M97935_MB_at
    (42447 total)
  varLabels: probeset gene EntrezGene.ID best_probe
  varMetadata: labelDescription
```

Details

assayData: 42447 features, 129 samples

Platform type:

Available sample meta-data:

sample_name:

Length	Class	Mode
129	character	character

alt_sample_name:

Length	Class	Mode
129	character	character

sample_type:

tumor

129

er:

negative positive

53 76

her2:

negative positive

98 31

tumor_size:

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
0.800	1.500	2.200	2.488	3.000	8.500

N:

0 1

64 65

grade:

1 2 3

27 32 70

treatment:

untreated

129

batch:

IRB

129

uncurated_author_metadata:

Length Class Mode
129 character character

Source

<http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE5460>

K00	KOO
-----	-----

Description

link does not work, in progress8

Format

```
experimentData(eset):
Experiment data
  Experimenter name:
  Laboratory:
  Contact information: http://www.ncbi.nlm.nih.gov/pubmed/12747878
  Title:
  URL: Unavailable
  PMIDs: 12747878

Abstract: A 6 word abstract is available. Use 'abstract' method.
notes:
  summary:
    A new gene signature was used to accurately predict 90
n the study.
  mapping.method:
    maxRowVariance
  mapping.group:
    EntrezGene.ID
  preprocessing:
    As published by original author.

featureData(eset):
An object of class 'AnnotatedDataFrame'
  featureNames: 1007_s_at 1053_at ... AFFX-HUMISGF3A/M97935_MB_at (280
    total)
  varLabels: probeset gene EntrezGene.ID best_probe
  varMetadata: labelDescription
```

Details

assayData: 280 features, 88 samples

Platform type:

Available sample meta-data:

sample_name:

Length	Class	Mode
88	character	character

alt_sample_name:

Length	Class	Mode
88	character	character

sample_type:

tumor
88

er:

negative	positive
15	73

pgr:

negative	positive
23	65

tumor_size:

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
0.20	1.80	2.30	2.74	3.50	8.50

N:

0	1
19	69

treatment:

chemotherapy	untreated
61	27

batch:

K00
88

uncurated_author_metadata:

Length	Class	Mode
88	character	character

duplicates:

Length	Class	Mode
88	character	character

Source

Unavailable

loadBreastDatasets	<i>Function to load breast cancer SummarizedExperiment objects from the Experiment Hub</i>
--------------------	--

Description

This function returns breast cancer datasets from the hub and a vector of patients from the datasets that are duplicates based on a spearman correlation > 0.98

Usage

```
loadBreastDatasets(  
  rescale = FALSE,  
  minNumberGenes = 0,  
  minNumberEvents = 0,  
  minSampleSize = 0,  
  keepCommonOnly = FALSE,  
  imputeMissing = FALSE,  
  removeDuplicates = FALSE  
)
```

Arguments

rescale	apply centering and scaling to the expression sets (default FALSE)
minNumberGenes	an integer specifying to remove expression sets with less genes than this number (default 0)
minNumberEvents	an integer specifying how man survival events must be in the dataset to keep the dataset (default 0)
minSampleSize	an integer specifying the minimum number of patients required in a summarizedExperiment (default 0)
keepCommonOnly	remove entrezIDs not common to all datasets (default FALSE)
imputeMissing	remove patients from datasets with missing expression values
removeDuplicates	remove patients with a Spearman correlation greater than or equal to 0.98 with other patient expression profiles (default TRUE)

Value

A 'list' with 2 elements. The First element named 'SummarizedExperiment's contains the datasets. The second element named duplicates contains a vector with patient IDs for the duplicate patients (those with Spearman correlation greater than or equal to 0.98 with other patient expression profiles).

loadBreastEsets	<i>Function to load breast cancer expression sets from the Experiment Hub</i>
-----------------	---

Description

This function returns breast cancer datasets from the hub and a vector of patients from the datasets that are most likely duplicates

Usage

```
loadBreastEsets(
  loadString = "majority",
  removeDuplicates = TRUE,
  quantileCutoff = 0,
  rescale = FALSE,
  minNumberGenes = 0,
  minNumberEvents = 0,
  minSampleSize = 0,
  removeRetracted = TRUE,
  removeSubsets = TRUE,
  keepCommonOnly = FALSE,
  imputeMissing = FALSE
)
```

Arguments

loadString	a character vector specifying which data will be loaded. The default is "majority", which loads in 37 of the 39 datasets. The other option is to provide a character vector of the names of the datasets to load. The metabric and tcga datasets are loaded separately as they are very large and doing so will help prevent memory allocation errors for R windows. Furthermore, these datasets are so large that they dominate statistical analyses so it is best that they are analyzed separate of the 37 smaller datasets loaded with the string majority
removeDuplicates	remove patients with a Spearman correlation greater than or equal to 0.98 with other patient expression profiles (default TRUE)
quantileCutoff	A numeric between 0 and 1 specifying to remove genes with standard deviation below the required quantile (default 0)
rescale	apply centering and scaling to the expression sets (default FALSE)

minNumberGenes an integer specifying to remove expression sets with less genes than this number (default 0)
minNumberEvents an integer specifying how many survival events must be in the dataset to keep the dataset (default 0)
minSampleSize an integer specifying the minimum number of patients required in an eset (default 0)
removeRetracted remove datasets from retracted papers (default TRUE, currently just PMID17290060 dataset)
removeSubsets remove datasets that are a subset of other datasets (default TRUE, currently just PMID19318476)
keepCommonOnly remove probes not common to all datasets (default FALSE)
imputeMissing remove patients from datasets with missing expression values

Value

a list with 2 elements. The first element named `esets` contains the datasets. The second element named `duplicates` contains a vector with patient IDs for the duplicate patients (those with Spearman correlation greater than or equal to 0.98 with other patient expression profiles).

Examples

```
## Use the default loadString="majority" if you want the 37 smaller datasets
esetsAndDups <- loadBreastEsets(loadString = c("CAL", "DFHCC", "DFHCC2",
  "DFHCC3", "DUKE", "DUKE2", "EMC2"))
```

LUND

LUND

Description

ExpressionSet for the LUND Dataset

Format

```

experimentData(eset):
Experiment data
  Experimenter name:
  Laboratory:
  Contact information: http://www.ncbi.nlm.nih.gov/pubmed/?term=18430221
  Title:
  URL: http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE31863
  PMIDs: 18430221
  No abstract available.
```

```

notes:
  summary:
    A significant difference was found between the ER positive subgroup and ER
    negative subgroup in the gene expression profiles.
  mapping.method:
    maxRowVariance
  mapping.group:
    EntrezGene.ID
  preprocessing:
    As published by original author.

featureData(eset):
An object of class 'AnnotatedDataFrame'
  featureNames: H200006618 H200006808 ... H300022925 (11154 total)
  varLabels: probeset gene EntrezGene.ID best_probe
  varMetadata: labelDescription

```

Details

```

assayData: 11154 features, 143 samples
Platform type:

```

```

-----
Available sample meta-data:
-----

```

```

sample_name:
  Length      Class      Mode
    143 character character

```

```

alt_sample_name:
  Length      Class      Mode
    143 character character

```

```

sample_type:
tumor
  143

```

```

er:
negative positive
   29      114

```

```

pgr:
negative positive  NA's
   47      88      8

```

```

tumor_size:
  Min. 1st Qu.  Median    Mean 3rd Qu.    Max.   NA's
  0.200  1.100  1.500  1.486  1.800  4.000     2

```

```

N:
  0
143

age_at_initial_pathologic_diagnosis:
  Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
  27.00  47.50   56.00   54.76  63.00   73.00

batch:
LUNDS1 LUNDS2 LUNDS3 LUNDS4
   30    47    22    44

uncurated_author_metadata:
  Length      Class      Mode
    143 character character

```

Source

<http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE31863>

LUND2

LUND2

Description

ExpressionSet for the LUND2 Dataset

Format

```

experimentData(eset):
Experiment data
  Experimenter name:
  Laboratory:
  Contact information: http://www.ncbi.nlm.nih.gov/pubmed/?term=17452630
  Title:
  URL: http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE5325
  PMIDs: 17452630
  No abstract available.
  notes:
    summary:
      Microarray signature was able to show PTEN mRNA losse_when IHC was unable,
      even though tumors exhibited PTEN loss behavior. Stathmim was an accurate
      IHC marker of the signature and had prognostic significance.
    mapping.method:
      maxRowVariance

```

```

mapping.group:
  EntrezGene.ID
preprocessing:
  As published by original author.

featureData(eset):
An object of class 'AnnotatedDataFrame'
  featureNames: 1 2 ... 27648 (22008 total)
  varLabels: probeset gene EntrezGene.ID best_probe
  varMetadata: labelDescription

```

Details

assayData: 22008 features, 105 samples

Platform type:

 Available sample meta-data:

sample_name:

Length	Class	Mode
105	character	character

alt_sample_name:

Length	Class	Mode
105	character	character

sample_type:

tumor
105

er:

negative	positive
60	45

treatment:

hormonotherapy
105

batch:

LUND2
105

uncurated_author_metadata:

Length	Class	Mode
105	character	character

Source

<http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE5325>

MAINZ

MAINZ

Description

ExpressionSet for the MAINZ Dataset

Format

```

experimentData(eset):
Experiment data
  Experimenter name:
  Laboratory:
  Contact information: http://www.ncbi.nlm.nih.gov/pubmed/?term=18593943
  Title:
  URL: http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE11121
  PMIDs: 18593943
  No abstract available.
  notes:
    summary:
      Poor prognosis is noted in tumors with low ER expression, showing the high
      est level of proliferative activity. In some tumors with highly expressed
      B-cell or T-cell metagenes, metastases rarely occurred, even with high pro
      liferation and low ER expression.
    mapping.method:
      maxRowVariance
    mapping.group:
      EntrezGene.ID
    preprocessing:
      As published by original author.

featureData(eset):
An object of class 'AnnotatedDataFrame'
  featureNames: 1007_s_at 1053_at ... AFFX-HUMISGF3A/M97935_MB_at
    (20967 total)
  varLabels: probeset gene EntrezGene.ID best_probe
  varMetadata: labelDescription

```

Details

assayData: 20967 features, 200 samples
Platform type:

Available sample meta-data:

sample_name:

Length	Class	Mode
200	character	character

alt_sample_name:

Length	Class	Mode
200	character	character

sample_type:

tumor
200

er:

negative	positive
38	162

tumor_size:

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
0.100	1.500	2.000	2.070	2.425	6.000

N:

0
200

age_at_initial_pathologic_diagnosis:

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
25.00	50.00	60.00	59.98	69.00	90.00

grade:

1	2	3
29	136	35

dmfs_days:

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
30	1905	2715	2816	3855	7200

dmfs_status:

norecurrence	recurrence
154	46

treatment:

untreated
200

batch:

MAINZ
200

uncurated_author_metadata:
Length Class Mode
200 character character

Source

<http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE11121>

MAQC2

MAQC2

Description

ExpressionSet for the MAQC2 Dataset

Format

```
experimentData(eset):
Experiment data
  Experimenter name:
  Laboratory:
  Contact information: http://www.ncbi.nlm.nih.gov/pubmed/?term=20064235
  Title:
  URL: http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE20194
  PMIDs: 20064235
  No abstract available.
  notes:
    summary:
      It is possible to build multi-gene classifiers of clinical outcome. Prediction accuracy depends on training sample size and classification difficulty.
  mapping.method:
    maxRowVariance
  mapping.group:
    EntrezGene.ID
  preprocessing:
    As published by original author.

featureData(eset):
An object of class 'AnnotatedDataFrame'
  featureNames: 1007_s_at 1053_at ... AFFX-HUMISGF3A/M97935_MB_at (20967 total)
  varLabels: probeset gene EntrezGene.ID best_probe
```

```
varMetadata: labelDescription
```

Details

```
assayData: 20967 features, 230 samples
```

```
Platform type:
```

```
-----  
Available sample meta-data:  
-----
```

```
sample_name:
```

Length	Class	Mode
230	character	character

```
alt_sample_name:
```

Length	Class	Mode
230	character	character

```
sample_type:
```

tumor
230

```
er:
```

negative	positive
89	141

```
pgr:
```

negative	positive
126	104

```
her2:
```

negative	positive
190	40

```
N:
```

0	1
66	164

```
age_at_initial_pathologic_diagnosis:
```

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
26.00	45.00	51.00	52.02	59.00	79.00

```
grade:
```

1	2	3
13	94	123

```
treatment:
```

```
chemotherapy
```

230

batch:

MAQC2

230

uncurated_author_metadata:

Length Class Mode

230 character character

Source<http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE20194>

MCCC

*MCCC***Description**

ExpressionSet for the MCCC Dataset

Format

experimentData(eset):

Experiment data

Experimenter name:

Laboratory:

Contact information: <http://www.ncbi.nlm.nih.gov/pubmed/?term=19960244>

Title:

URL: <http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE19177>

PMIDs: 19960244

No abstract available.

notes:

summary:

Overall, expression and copy number profiling of familial tumors have shown that the tumors show molecular heterogeneity similar to sporadic tumors and are defined by their molecular subtypes rather than BRCA1 or BRCA2 germline mutation status.

mapping.method:

maxRowVariance

mapping.group:

EntrezGene.ID

preprocessing:

As published by original author.

featureData(eset):

```
An object of class 'AnnotatedDataFrame'
  featureNames: probe_10017 probe_10021 ... probe_7650767 (19048 total)
  varLabels: probeset gene EntrezGene.ID best_probe
  varMetadata: labelDescription
```

Details

```
assayData: 19048 features, 75 samples
Platform type:
```

```
-----
Available sample meta-data:
-----
```

```
sample_name:
  Length      Class      Mode
      75 character character
```

```
sample_type:
tumor
      75
```

```
batch:
MCCC
      75
```

```
uncurated_author_metadata:
  Length      Class      Mode
      75 character character
```

Source

<http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE19177>

MDA4

MDA4

Description

ExpressionSet for the MDA4 Dataset

Format

```
experimentData(eset):
Experiment data
  Experimenter name:
```

```

Laboratory:
Contact information: http://www.ncbi.nlm.nih.gov/pubmed/?term=16896004
Title:
URL: http://bioinformatics.mdanderson.org/pubdata.html
PMIDs: 16896004
No abstract available.
notes:
  summary:
    The developed 30-probe set has high sensitivity and negative predictive value, accurately identifying 12 out of 13 patients with pCR and 27 out of 28 patients with residual disease.
  mapping.method:
    maxRowVariance
  mapping.group:
    EntrezGene.ID
  preprocessing:
    As published by original author.

featureData(eset):
An object of class 'AnnotatedDataFrame'
  featureNames: 1007_s_at 1053_at ... AFFX-HUMISGF3A/M97935_MB_at
  (21169 total)
  varLabels: probeset gene EntrezGene.ID best_probe
  varMetadata: labelDescription

```

Details

```

assayData: 21169 features, 129 samples
Platform type:
-----

```

```

Available sample meta-data:
-----

```

```

sample_name:
  Length      Class      Mode
    129 character character

```

```

unique_patient_ID:
  Length      Class      Mode
    129 character character

```

```

sample_type:
tumor
  129

```

```

er:
negative positive  NA's
    48      79      2

```

pgr:
negative positive NA's
73 54 2

her2:
negative positive
114 15

tumor_size:
Min. 1st Qu. Median Mean 3rd Qu. Max. NA's
0.000 0.500 1.800 2.162 3.000 10.000 8

N:
0 1 NA's
59 62 8

age_at_initial_pathologic_diagnosis:
Min. 1st Qu. Median Mean 3rd Qu. Max.
28.00 43.00 51.00 51.43 61.00 73.00

treatment:
chemotherapy
129

batch:
MDA4
129

uncurated_author_metadata:
Length Class Mode
129 character character

duplicates:
MDA4.MDA4_M207 MDA4.MDA4_M400 NA's
1 1 127

Source

<http://bioinformatics.mdanderson.org/pubdata.html>

METABRIC	METABRIC
----------	----------

Description

ExpressionSet for the METABRIC Dataset

Format

```

experimentData(eset):
Experiment data
  Experimenter name:
  Laboratory:
  Contact information: http://www.ncbi.nlm.nih.gov/pubmed/22522925
  Title:
  URL: https://www.ebi.ac.uk/ega/studies/EGAS00000000083
  PMIDs: 22522925
  No abstract available.
notes:
  summary:

  mapping.method:
    maxRowVariance
  mapping.group:
    EntrezGene.ID
  preprocessing:
    As published by original author.

featureData(eset):
An object of class 'AnnotatedDataFrame'
  featureNames: ILMN_1802380 ILMN_1736104 ... ILMN_1709472 (36155
    total)
  varLabels: probeset gene EntrezGene.ID best_probe
  varMetadata: labelDescription

```

Details

```

assayData: 36155 features, 2136 samples
Platform type:
Overall survival time-to-event summary (in years):
Call: survfit(formula = Surv(time, cens) ~ -1)

```

```

      165 observations deleted due to missingness
      n  events  median 0.95LCL 0.95UCL
1971.0   891.0   12.3    11.6    13.2

```

```

-----
Available sample meta-data:
-----

```

```

sample_name:
  Length      Class      Mode
    2136 character character

```

```

alt_sample_name:

```

```

      Length      Class      Mode
      2136 character character

sample_type:
healthy  tumor
   144    1992

er:
negative positive  NA's
   440    1508    188

her2:
negative positive  NA's
   676    148    1312

tumor_size:
  Min. 1st Qu.  Median    Mean 3rd Qu.    Max.   NA's
  0.000   1.700   2.300   2.621   3.000  18.200   164

N:
  0    1 NA's
1042  950 144

age_at_initial_pathologic_diagnosis:
  Min. 1st Qu.  Median    Mean 3rd Qu.    Max.   NA's
 21.93   51.36   61.78   61.13   70.76   96.29    13

grade:
  1    2    3 NA's
 170  775  957  234

days_to_death:
  Min. 1st Qu.  Median    Mean 3rd Qu.    Max.   NA's
    3    1498   2632   2948   4357   9218   147

vital_status:
deceased  living   NA's
   891    1081    164

treatment:
chemo.plus.hormono  chemotherapy  hormonotherapy  untreated
           196           226           1029           685

batch:
  Min. 1st Qu.  Median    Mean 3rd Qu.    Max.   NA's
 1.000   1.000   3.000   2.613   3.000   5.000   144

uncurated_author_metadata:

```

```

      Length      Class      Mode
      2136 character character

duplicates:
      Length      Class      Mode
      2136 character character
```

Source

<https://www.ebi.ac.uk/ega/studies/EGAS000000000083>

MSK	<i>MSK</i>
-----	------------

Description

ExpressionSet for the MSK Dataset

Format

```

experimentData(eset):
Experiment data
  Experimenter name:
  Laboratory:
  Contact information: http://www.ncbi.nlm.nih.gov/pubmed/?term=16049480
  Title:
  URL: http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=gse2603
  PMIDs: 16049480
  No abstract available.
notes:
  summary:
    A set of genes were identified that mark and mediate metastasis to the lung.
    Some genes confer growth advantages to both the breast tumor and lung environment,
    while others contribute to aggressive growth specifically in the lung.
  mapping.method:
    maxRowVariance
  mapping.group:
    EntrezGene.ID
  preprocessing:
    As published by original author.

featureData(eset):
An object of class 'AnnotatedDataFrame'
  featureNames: 1007_s_at 1053_at ... AFFX-HUMISGF3A/M97935_MB_at
  (20967 total)
```

```
varLabels: probeset gene EntrezGene.ID best_probe
varMetadata: labelDescription
```

Details

```
assayData: 20967 features, 99 samples
Platform type:
```

```
-----
Available sample meta-data:
-----
```

```
sample_name:
  Length      Class      Mode
    99 character character
```

```
alt_sample_name:
  Length      Class      Mode
    99 character character
```

```
sample_type:
tumor
  99
```

```
er:
negative positive
   42         57
```

```
pgr:
negative positive  NA's
   55         43     1
```

```
her2:
positive  NA's
   85     14
```

```
tumor_size:
  Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
  1.100  2.450   3.200   3.624  4.300  10.000
```

```
N:
  0  1
34 65
```

```
age_at_initial_pathologic_diagnosis:
  Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
  30.00  46.50   56.00   55.81  63.50   87.00
```

```
dmfs_days:
```

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.	NA's
245	1279	1971	1888	2575	3924	17
dmfs_status:						
norecurrence	recurrence		NA's			
55	27		17			
batch:						
MSK						
99						
uncurated_author_metadata:						
Length		Class	Mode			
99		character	character			

Source

<http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=gse2603>

MUG	MUG
-----	-----

Description

ExpressionSet for the MUG Dataset

Format

```
experimentData(eset):
Experiment data
  Experimenter name:
  Laboratory:
  Contact information: http://www.ncbi.nlm.nih.gov/pubmed/?term=18592372
  Title:
  URL: http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=gse10510
  PMIDs: 18592372
  No abstract available.
notes:
  summary:
    A method was developed to separate tumor cells and their microenvironment
    to test the prognostic abilities of the immune system. Results showed that
    lymphatic infiltration is beneficial for ER negative patients, but probab
    ly not beneficial for ER positive patients.
  mapping.method:
    maxRowVariance
  mapping.group:
```

```

      EntrezGene.ID
preprocessing:
  As published by original author.

featureData(eset):
An object of class 'AnnotatedDataFrame'
  featureNames: H200000001 H200000005 ... opHsV04TC000043 (14288 total)
  varLabels: probeset gene EntrezGene.ID best_probe
  varMetadata: labelDescription

```

Details

```

assayData: 14288 features, 152 samples
Platform type:
-----

```

```

Available sample meta-data:
-----

```

```

sample_name:
  Length      Class      Mode
    152 character character

```

```

alt_sample_name:
  Length      Class      Mode
    152 character character

```

```

sample_type:
tumor
  152

```

```

batch:
MUG
  152

```

```

uncurated_author_metadata:
  Length      Class      Mode
    152 character character

```

Source

<http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=gse10510>

Description

ExpressionSet for the NCCS Dataset

Format

```

experimentData(eset):
Experiment data
  Experimenter name:
  Laboratory:
  Contact information: http://www.ncbi.nlm.nih.gov/pubmed/?term=18636107
  Title:
  URL: http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=gse5364
  PMIDs: 18636107
  No abstract available.
  notes:
    summary:
      48 genes were identified that displayed highly restricted levels of expres
sion in tumors compared to normal tissues. This was validated in 11 indepe
ndent cohorts of different cancer types.
    mapping.method:
      maxRowVariance
    mapping.group:
      EntrezGene.ID
    preprocessing:
      As published by original author.

featureData(eset):
An object of class 'AnnotatedDataFrame'
  featureNames: 1007_s_at 1053_at ... AFFX-HUMISGF3A/M97935_MB_at
  (20967 total)
  varLabels: probeset gene EntrezGene.ID best_probe
  varMetadata: labelDescription

```

Details

```

assayData: 20967 features, 183 samples
Platform type:
-----
Available sample meta-data:
-----

sample_name:
  Length      Class      Mode
    183 character character

alt_sample_name:
  Min. 1st Qu.  Median    Mean 3rd Qu.    Max.

```

1.0 46.5 92.0 92.0 137.5 183.0

sample_type:

tumor

183

batch:

NCCS

183

uncurated_author_metadata:

Length Class Mode

183 character character

Source

<http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=gse5364>

NCI

NCI

Description

ExpressionSet for the NCI Dataset

Format

experimentData(eset):

Experiment data

Experimenter name:

Laboratory:

Contact information: <http://www.ncbi.nlm.nih.gov/pubmed/?term=12917485>

Title:

URL: Supplemental data from paper

PMIDs: 12917485

No abstract available.

notes:

summary:

Expression patterns were strongly associated with ER status, moderately associated with grade, but not associated with menopausal state, node status, or tumor size. Genes that were significantly associated with survival were identified.

mapping.method:

maxRowVariance

mapping.group:

EntrezGene.ID

```
preprocessing:
  As published by original author.
```

```
featureData(eset):
An object of class 'AnnotatedDataFrame'
  featureNames: AF106966 AF217974 ... Y12473 (5154 total)
  varLabels: probeset gene EntrezGene.ID best_probe
  varMetadata: labelDescription
```

Details

```
assayData: 5154 features, 99 samples
Platform type:
```

```
-----
Available sample meta-data:
-----
```

```
sample_name:
  Length      Class      Mode
    99 character character
```

```
alt_sample_name:
  Min. 1st Qu.  Median    Mean 3rd Qu.  Max.
 21580  21610   21640   21650  21670   21830
```

```
sample_type:
tumor
  99
```

```
er:
negative positive
   34      65
```

```
tumor_size:
  Min. 1st Qu.  Median    Mean 3rd Qu.  Max.
  0.80   1.80   2.50   2.82   3.00   8.00
```

```
N:
  0  1
46 53
```

```
age_at_initial_pathologic_diagnosis:
  Min. 1st Qu.  Median    Mean 3rd Qu.  Max.
 33.00  49.00  57.00  57.47  64.50  90.00
```

```
grade:
  1  2  3
16 38 45
```

```

days_to_tumor_recurrence:
  Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
      8     967     2057    1969    2930    4067

```

```

recurrence_status:
norecurrence  recurrence
           54           45

```

```

treatment:
  chemotherapy  hormonotherapy  untreated
           10           78           11

```

```

batch:
NCI
99

```

```

uncurated_author_metadata:
  Length      Class      Mode
    99 character character

```

Source

Supplemental data from paper

NKI

NKI

Description

ExpressionSet for the NKI Dataset

Format

```

experimentData(eset):
Experiment data
  Experimenter name:
  Laboratory:
Contact information: http://www.ncbi.nlm.nih.gov/pubmed/?term=12490681; http://www.ncbi.nlm.nih.gov
Title:
URL: Not available
PMIDs: 12490681, 11823860
No abstract available.
notes:
  summary:
    It was found that the gene expression profile that was studied was more po

```

werful in predicting outcome of disease in younger patients than using standard clinical and pathological criteria.

```
mapping.method:
  maxRowVariance
mapping.group:
  EntrezGene.ID
preprocessing:
  As published by original author.
```

```
featureData(eset):
```

```
An object of class 'AnnotatedDataFrame'
```

```
featureNames: Contig45645_RC Contig44916_RC ... Contig62037_RC (14960
total)
varLabels: probeset gene EntrezGene.ID best_probe
varMetadata: labelDescription
```

Details

```
assayData: 14960 features, 337 samples
```

```
Platform type:
```

```
Overall survival time-to-event summary (in years):
```

```
Call: survfit(formula = Surv(time, cens) ~ -1)
```

```
42 observations deleted due to missingness
```

n	events	median	0.95LCL	0.95UCL
295	79	NA	NA	NA

```
-----
Available sample meta-data:
-----
```

```
sample_name:
```

Length	Class	Mode
337	character	character

```
alt_sample_name:
```

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
4.0	123.0	215.0	214.1	312.0	404.0

```
sample_type:
```

```
tumor
```

```
337
```

```
er:
```

```
negative positive
```

```
88 249
```

```
tumor_size:
```

	Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
	0.200	1.500	2.000	2.241	2.800	5.500

N:

0	1
193	144

age_at_initial_pathologic_diagnosis:

	Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
	26.0	40.0	45.0	44.2	49.0	62.0

grade:

1	2	3
79	109	149

dmfs_days:

	Min.	1st Qu.	Median	Mean	3rd Qu.	Max.	NA's
	9	1252	2414	2546	3602	6699	18

dmfs_status:

norecurrence	recurrence	NA's
210	109	18

days_to_tumor_recurrence:

	Min.	1st Qu.	Median	Mean	3rd Qu.	Max.	NA's
	9	1252	2414	2546	3602	6699	18

recurrence_status:

norecurrence	recurrence	NA's
210	109	18

days_to_death:

	Min.	1st Qu.	Median	Mean	3rd Qu.	Max.	NA's
	20	1934	2637	2870	3763	6694	42

vital_status:

deceased	living	NA's
79	216	42

treatment:

chemotherapy	hormonotherapy	untreated
90	40	207

batch:

NKI	NKI2
117	220

uncurated_author_metadata:

Length Class Mode
337 character character

Source

Not available

PNC	<i>PNC</i>
-----	------------

Description

ExpressionSet for the PNC Dataset

Format

```
experimentData(eset):
Experiment data
  Experimenter name:
  Laboratory:
  Contact information: http://www.ncbi.nlm.nih.gov/pubmed/?term=21910250
  Title:
  URL: http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE20711
  PMIDs: GSE20711, PMID 21910250
  No abstract available.
  notes:
    summary:
      Breast tumors can be further divided than the currently known expression s
ubtypes based on DNA methylation profiles.
    mapping.method:
      maxRowVariance
    mapping.group:
      EntrezGene.ID
    preprocessing:
      As published by original author.

featureData(eset):
An object of class 'AnnotatedDataFrame'
  featureNames: 1007_s_at 1053_at ... AFFX-HUMISGF3A/M97935_MB_at
  (42447 total)
  varLabels: probeset gene EntrezGene.ID best_probe
  varMetadata: labelDescription
```

Details

assayData: 42447 features, 92 samples
 Platform type:
 Overall survival time-to-event summary (in years):
 Call: survfit(formula = Surv(time, cens) ~ -1)

4 observations deleted due to missingness

n	events	median	0.95LCL	0.95UCL
88.0	25.0	NA	11.3	NA

 Available sample meta-data:

sample_name:

Length	Class	Mode
92	character	character

alt_sample_name:

Length	Class	Mode
92	character	character

sample_type:

tumor
92

er:

negative	positive	NA's
43	45	4

pgr:

negative	positive	NA's
43	40	9

her2:

negative	positive	NA's
64	26	2

tumor_size:

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.	NA's
0.900	1.700	2.500	2.758	3.000	10.000	6

N:

0	1	NA's
43	40	9

age_at_initial_pathologic_diagnosis:

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.	NA's
------	---------	--------	------	---------	------	------

```

32.16  48.57  53.90  55.97  64.84  82.13      4

grade:
  1    2    3 NA's
13    5   70    4

days_to_tumor_recurrence:
  Min. 1st Qu.  Median    Mean 3rd Qu.    Max.    NA's
    29    967   2216   2122   2931   5139     7

recurrence_status:
norecurrence  recurrence      NA's
          49           36          7

days_to_death:
  Min. 1st Qu.  Median    Mean 3rd Qu.    Max.    NA's
   318   1940   2372   2525   3043   5139     4

vital_status:
deceased  living    NA's
    25      63      4

batch:
PNC
92

uncurated_author_metadata:
  Length    Class      Mode
    92 character character
```

Source

<http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE20711>

STK	STK
-----	-----

Description

ExpressionSet for the STK Dataset

Format

```
experimentData(eset):
Experiment data
  Experimenter name:
```

```

Laboratory:
Contact information: http://www.ncbi.nlm.nih.gov/pubmed/?term=16280042
Title:
URL: http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=gse1456
PMIDs: 16280042
No abstract available.
notes:
  summary:
    Expression profiling was able to better predict prognosis compared to histological staging.
  mapping.method:
    maxRowVariance
  mapping.group:
    EntrezGene.ID
  preprocessing:
    As published by original author.

featureData(eset):
An object of class 'AnnotatedDataFrame'
  featureNames: 1007_s_at 1053_at ... 244889_at (36178 total)
  varLabels: probeset gene EntrezGene.ID best_probe
  varMetadata: labelDescription

```

Details

```

assayData: 36178 features, 159 samples
Platform type:
-----
Available sample meta-data:
-----

sample_name:
  Length      Class      Mode
    159 character character

alt_sample_name:
  Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
   1.0   67.0   136.0   138.3  208.5   277.0

sample_type:
tumor
  159

er:
negative positive
   29      130

age_at_initial_pathologic_diagnosis:

```

```

      Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
      31.0   48.0   56.0   57.8   68.5   87.0

grade:
  1     2     3 NA's
28   58   61   12

days_to_tumor_recurrence:
      Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
      66   2022   2467   2234   2846   3099

recurrence_status:
norecurrence  recurrence
      113           46

treatment:
  chemotherapy hormonotherapy      untreated
           89           48           22

batch:
STK
159

uncurated_author_metadata:
      Length      Class      Mode
      159 character character

```

Source

<http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=gse1456>

STNO2

STNO2

Description

ExpressionSet for the STNO2 Dataset

Format

```

experimentData(eset):
Experiment data
  Experimenter name:
  Laboratory:
  Contact information: http://www.ncbi.nlm.nih.gov/pubmed/?term=12829800
  Title:

```

```

URL: http://smd.princeton.edu/cgi-bin/publication/viewPublication.pl?pub_no=248
PMIDs: 12829800
No abstract available.
notes:
  summary:
    Distinct breast cancer subtypes were determined by gene expression profile
s and were validated in other published datasets.
  mapping.method:
    maxRowVariance
  mapping.group:
    EntrezGene.ID
  preprocessing:
    As published by original author.

featureData(eset):
An object of class 'AnnotatedDataFrame'
  featureNames: IMAGE:1020315 IMAGE:1030271 ... IMAGE:971399 (3663
    total)
  varLabels: probeset gene EntrezGene.ID best_probe
  varMetadata: labelDescription

```

Details

```

assayData: 3663 features, 118 samples
Platform type:
Overall survival time-to-event summary (in years):
Call: survfit(formula = Surv(time, cens) ~ -1)

```

n	events	median	0.95LCL	0.95UCL
118.00	46.00	4.67	3.34	NA

```

-----
Available sample meta-data:
-----

```

```

sample_name:
  Length      Class      Mode
    118 character character

```

```

alt_sample_name:
  Length      Class      Mode
    118 character character

```

```

sample_type:
tumor
  118

```

```

er:

```

negative	positive	NA's
31	82	5

tumor_size:				
1	2	3	4	NA's
6	13	62	32	5

N:		
0	1	NA's
34	79	5

age_at_initial_pathologic_diagnosis:					
Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
21.00	46.25	58.00	58.47	71.75	85.00

grade:			
1	2	3	NA's
11	49	53	5

days_to_tumor_recurrence:						
Min.	1st Qu.	Median	Mean	3rd Qu.	Max.	NA's
122.0	396.0	761.0	927.9	1233.0	2800.0	23

recurrence_status:	
norecurrence	recurrence
58	60

days_to_death:					
Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
91	426	898	1019	1392	5722

vital_status:	
deceased	living
46	72

treatment:		
chemotherapy	hormonotherapy	untreated
23	73	22

batch:
STNO2
118

uncurated_author_metadata:		
Length	Class	Mode
118	character	character

Source

http://smd.princeton.edu/cgi-bin/publication/viewPublication.pl?pub_no=248

TCGA

TCGA

Description

ExpressionSet for the TCGA Dataset

Format

```

experimentData(eset):
Experiment data
  Experimenter name:
  Laboratory:
  Contact information: http://www.ncbi.nlm.nih.gov/pubmed/23000897
  Title:
  URL: http://cancergenome.nih.gov/
  PMIDs: 23000897
  No abstract available.
  notes:
    summary:

  mapping.method:
    maxRowVariance
  mapping.group:
    EntrezGene.ID
  preprocessing:
    As published by original author.

featureData(eset):
An object of class 'AnnotatedDataFrame'
  featureNames: A1BG A2M ... ARHGAP11A.2 (19504 total)
  varLabels: probeset gene EntrezGene.ID best_probe
  varMetadata: labelDescription

```

Details

```

assayData: 19504 features, 1073 samples
Platform type:
Overall survival time-to-event summary (in years):
Call: survfit(formula = Surv(time, cens) ~ -1)

```

	n	events	median	0.95LCL	0.95UCL
	1073.00	103.00	10.05	8.56	12.05

```

-----
Available sample meta-data:
-----

sample_name:
  Length   Class      Mode
    1073 character character

alt_sample_name:
  Length   Class      Mode
    1073 character character

unique_patient_ID:
  Length   Class      Mode
    1073 character character

sample_type:
tumor
  1073

er:
negative positive  NA's
   233      790    50

pgr:
negative positive  NA's
   334      686    53

her2:
negative positive  NA's
   549      161   363

age_at_initial_pathologic_diagnosis:
  Min. 1st Qu.  Median    Mean 3rd Qu.  Max.
  26.00  49.00   58.00   58.48  68.00   90.00

days_to_death:
  Min. 1st Qu.  Median    Mean 3rd Qu.  Max.
   -7.0   137.0   412.0   820.3 1180.0 6796.0

vital_status:
deceased  living
    103     970

batch:
TCGA
1073

```

```

uncurated_author_metadata:
  Length      Class      Mode
    1073 character character

```

Source

<http://cancergenome.nih.gov/>

TRANSBIG

TRANSBIG

Description

ExpressionSet for the TRANSBIG Dataset

Format

```

experimentData(eset):
Experiment data
  Experimenter name:
  Laboratory:
  Contact information: http://www.ncbi.nlm.nih.gov/pubmed/?term=17545524
  Title:
  URL: http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=gsE7390
  PMIDs: 17545524
  No abstract available.
  notes:
    summary:
      The 76-gene signature was validated. The results supports the hypothesis t
      hat utilizing the gene signature could reduce the number of patients who r
      eceive unnecessary adjuvant therapy.
    mapping.method:
      maxRowVariance
    mapping.group:
      EntrezGene.ID
    preprocessing:
      As published by original author.

featureData(eset):
An object of class 'AnnotatedDataFrame'
  featureNames: 1007_s_at 1053_at ... AFFX-HUMISGF3A/M97935_MB_at
    (20967 total)
  varLabels: probeset gene EntrezGene.ID best_probe
  varMetadata: labelDescription

```

Details

assayData: 20967 features, 198 samples

Platform type:

Overall survival time-to-event summary (in years):

Call: survfit(formula = Surv(time, cens) ~ -1)

n	events	median	0.95LCL	0.95UCL
198.0	56.0	NA	17.1	NA

Available sample meta-data:

sample_name:

Length	Class	Mode
198	character	character

sample_type:

tumor
198

er:

negative	positive
64	134

tumor_size:

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
0.600	1.700	2.000	2.181	2.500	5.000

N:

0
198

age_at_initial_pathologic_diagnosis:

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
24.00	42.00	46.00	46.39	51.00	60.00

grade:

1	2	3	NA's
30	83	83	2

dmfs_days:

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
125	2375	4384	3954	5566	9108

dmfs_status:

norecurrence	recurrence
147	51

```

days_to_tumor_recurrence:
  Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
    121    1528    3534    3399    5130    8711

recurrence_status:
norecurrence  recurrence
         112             86

days_to_death:
  Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
    146    2744    4562    4150    5610    9108

vital_status:
deceased  living
         56     142

treatment:
untreated
        198

batch:
VDXGUYU VDXIGRU VDXKIU VDXOXFU VDXRHU
      36      50      51      24      37

uncurated_author_metadata:
  Length    Class      Mode
    198 character character

```

Source

<http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=gsE7390>

UCSF

UCSF

Description

ExpressionSet for the UCSF Dataset

Format

```

experimentData(eset):
Experiment data
  Experimenter name:
  Laboratory:

```

Contact information: <http://www.ncbi.nlm.nih.gov/pubmed/?term=17428335>; <http://www.ncbi.nlm.nih.gov>

Title:

URL: Not available

PMIDs: 17428335, 14612510

No abstract available.

notes:

summary:

A gene set was identified that correctly predicted outcomes more effectively than using histological markers.

mapping.method:

maxRowVariance

mapping.group:

EntrezGene.ID

preprocessing:

As published by original author.

featureData(eset):

An object of class 'AnnotatedDataFrame'

featureNames: probe_1 probe_3 ... probe_10365 (8015 total)

varLabels: probeset gene EntrezGene.ID best_probe

varMetadata: labelDescription

Details

assayData: 8015 features, 162 samples

Platform type:

Overall survival time-to-event summary (in years):

Call: survfit(formula = Surv(time, cens) ~ -1)

29 observations deleted due to missingness

n	events	median	0.95LCL	0.95UCL
133.00	44.00	11.56	9.25	NA

Available sample meta-data:

sample_name:

Length	Class	Mode
162	character	character

alt_sample_name:

Length	Class	Mode
162	character	character

sample_type:

tumor

162

```

er:
negative positive  NA's
    41      101      20

pgr:
negative positive  NA's
    46      94      22

her2:
negative positive  NA's
    35      19      108

tumor_size:
  Min. 1st Qu.  Median    Mean 3rd Qu.  Max.   NA's
  0.000  1.800  2.000  2.682  3.200  11.000     7

N:
  0      1 NA's
 67    82   13

age_at_initial_pathologic_diagnosis:
  Min. 1st Qu.  Median    Mean 3rd Qu.  Max.   NA's
 28.00  44.00  53.00  56.61  70.00  88.00     9

grade:
  1      2      3 NA's
 14    62    74   12

dmfs_days:
  Min. 1st Qu.  Median    Mean 3rd Qu.  Max.   NA's
   47     897   2040   2084   2992   8267     29

dmfs_status:
norecurrence  recurrence
      140           22

days_to_tumor_recurrence:
  Min. 1st Qu.  Median    Mean 3rd Qu.  Max.   NA's
   47     861   1865   1985   2847   8267     29

recurrence_status:
norecurrence  recurrence
      125           37

days_to_death:
  Min. 1st Qu.  Median    Mean 3rd Qu.  Max.   NA's
   47     1087   2054   2140   3087   8267     29

```

```
vital_status:
deceased    living    NA's
      54         99         9

treatment:
chemo.plus.hormono      chemotherapy      hormonotherapy      untreated
           31              38              61              22
           NA's
           10

batch:
UCSF
162

uncurated_author_metadata:
  Length      Class      Mode
    162 character character
```

Source

Not available

UNC4	UNC4
------	------

Description

ExpressionSet for the UNC4 Dataset

Format

```
experimentData(eset):
Experiment data
  Experimenter name:
  Laboratory:
  Contact information: http://www.ncbi.nlm.nih.gov/pubmed/?term=20813035
  Title:
  URL: http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=gse18229
  PMIDs: 20813035
  No abstract available.
notes:
  summary:
    Clinically, this subtype is usually triple negative invasive ductal carcinomas with a poor prognosis. Response to standard of care preoperative chemotherapy is intermediate between basal-like and luminal tumors. The claudin
```

n-low subtype most closely resembles the mammary epithelial stem cell.

mapping.method:
maxRowVariance

mapping.group:
EntrezGene.ID

preprocessing:
As published by original author.

featureData(eset):

An object of class 'AnnotatedDataFrame'

featureNames: probe.10 probe.12 ... probe.79701 (5420 total)

varLabels: probeset gene EntrezGene.ID best_probe

varMetadata: labelDescription

Details

assayData: 5420 features, 305 samples

Platform type:

Overall survival time-to-event summary (in years):

Call: survfit(formula = Surv(time, cens) ~ -1)

65 observations deleted due to missingness

n	events	median	0.95LCL	0.95UCL
240.00	51.00	7.73	6.82	NA

Available sample meta-data:

sample_name:

Length	Class	Mode
305	character	character

sample_type:

tumor

305

er:

negative	positive	NA's
99	154	52

pgr:

negative	positive	NA's
126	109	70

her2:

negative	positive	NA's
203	58	44

```

tumor_size:
  1  1.5  3  6 NA's
60  1 129 43 72

N:
  0  1 NA's
126 135 44

age_at_initial_pathologic_diagnosis:
  Min. 1st Qu.  Median    Mean 3rd Qu.    Max.    NA's
 24.00  46.00  55.00  56.73  68.00  89.00    59

grade:
  1  2  3 NA's
25 80 138 62

days_to_tumor_recurrence:
  Min. 1st Qu.  Median    Mean 3rd Qu.    Max.    NA's
 30.0  450.0  750.0  954.3 1380.0 3540.0    64

recurrence_status:
norecurrence  recurrence    NA's
          170           70         65

days_to_death:
  Min. 1st Qu.  Median    Mean 3rd Qu.    Max.    NA's
   30    540    885   1104   1590   5190    65

vital_status:
deceased  living    NA's
    51    189    65

batch:
UNC4
305

uncurated_author_metadata:
  Length    Class    Mode
    305 character character

duplicates:
K00.K00_KF_105 K00.K00_T01_514    NA's
              1              1    303

```

Source

<http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=gse18229>

UNT

UNT

Description

ExpressionSet for the UNT Dataset

Format

```
experimentData(eset):
```

```
Experiment data
```

```
  Experimenter name:
```

```
  Laboratory:
```

```
  Contact information: http://www.ncbi.nlm.nih.gov/pubmed/?term=16478745; http://www.ncbi.nlm.nih.gov
```

```
  Title:
```

```
  URL: http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=gse2990
```

```
  PMIDs: 16478745, 17401012
```

```
  No abstract available.
```

```
  notes:
```

```
    summary:
```

```
      A gene expression grading index (GGI) was developed. The GGI reclassified
      grade 2 patients into two groups with low and high risks of recurrence.
```

```
    mapping.method:
```

```
      maxRowVariance
```

```
    mapping.group:
```

```
      EntrezGene.ID
```

```
    preprocessing:
```

```
      As published by original author.
```

```
featureData(eset):
```

```
An object of class 'AnnotatedDataFrame'
```

```
  featureNames: 1007_s_at 1053_at ... 244889_at (36084 total)
```

```
  varLabels: probeset gene EntrezGene.ID best_probe
```

```
  varMetadata: labelDescription
```

Details

```
assayData: 36084 features, 133 samples
```

```
Platform type:
```

```
-----
```

```
Available sample meta-data:
```

```
-----
```

```
sample_name:
```

```
  Length      Class      Mode
```

```
    133 character character
```

```

alt_sample_name:
  Length      Class      Mode
    133 character character

sample_type:
tumor
  133

er:
negative positive  NA's
   40      86      7

pgr:
negative positive  NA's
   6      56      71

tumor_size:
  Min. 1st Qu.  Median    Mean 3rd Qu.  Max.
  0.000  1.200  1.900  1.892  2.300  6.000

N:
  0
133

age_at_initial_pathologic_diagnosis:
  Min. 1st Qu.  Median    Mean 3rd Qu.  Max.
  24.00  44.00  53.00  51.79  60.00  73.00

grade:
  1    2    3 NA's
  32   51   29   21

dmfs_days:
  Min. 1st Qu.  Median    Mean 3rd Qu.  Max.
   61   1338   2809   2724   4078   5305

dmfs_status:
norecurrence  recurrence  NA's
          97          28          8

days_to_tumor_recurrence:
  Min. 1st Qu.  Median    Mean 3rd Qu.  Max.
   61   1338   2675   2687   3912   5305

recurrence_status:
norecurrence  recurrence  NA's
          76          49          8

```

```

treatment:
untreated
    133

batch:
    KIU OXFU
    64  69

uncurated_author_metadata:
    Length      Class      Mode
    133 character character

```

Source

<http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=gse2990>

UPP	<i>UPP</i>
-----	------------

Description

ExpressionSet for the UPP Dataset

Format

```

experimentData(eset):
Experiment data
  Experimenter name:
  Laboratory:
  Contact information: http://www.ncbi.nlm.nih.gov/pubmed/?term=16141321
  Title:
  URL: http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=gse3494
  PMIDs: 16141321
  No abstract available.
  notes:
    summary:
      A 32-gene expression signature of p53 was identified that differentiates p
      -53 mutant and wild-type tumors. The signature is more effective than sequ
      ence-based assessments of p53 in predicting prognosis and therapeutic resp
      onse.
    mapping.method:
      maxRowVariance
    mapping.group:
      EntrezGene.ID
    preprocessing:

```

As published by original author.

```
featureData(eset):
An object of class 'AnnotatedDataFrame'
  featureNames: 1007_s_at 1053_at ... 244889_at (36178 total)
  varLabels: probeset gene EntrezGene.ID best_probe
  varMetadata: labelDescription
```

Details

```
assayData: 36178 features, 251 samples
Platform type:
```

```
-----
Available sample meta-data:
-----
```

```
sample_name:
  Length      Class      Mode
    251 character character
```

```
alt_sample_name:
  Length      Class      Mode
    251 character character
```

```
sample_type:
tumor
  251
```

```
er:
negative positive  NA's
    34      213      4
```

```
pgr:
negative positive
    61      190
```

```
tumor_size:
  Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
  0.200  1.500   2.000   2.243  2.562  13.000
```

```
N:
  0      1 NA's
158  84      9
```

```
age_at_initial_pathologic_diagnosis:
  Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
 28.00  52.00   64.00   62.11  72.00   93.00
```

```
grade:
  1      2      3 NA's
67 128  54      2

days_to_tumor_recurrence:
  Min. 1st Qu.  Median    Mean 3rd Qu.    Max.    NA's
   30   1870   3711   3007   3985   4654     17

recurrence_status:
norecurrence  recurrence      NA's
         181          55          15

treatment:
hormonotherapy      untreated      NA's
         80          142          29

batch:
UPPT UPPU
  80 171

uncurated_author_metadata:
  Length      Class      Mode
    251 character character
```

Source

<http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=gse3494>

VDX	VDX
-----	-----

Description

ExpressionSet for the VDX Dataset

Format

```
experimentData(eset):
Experiment data
  Experimenter name:
  Laboratory:
Contact information: http://www.ncbi.nlm.nih.gov/pubmed/?term=15721472; http://www.ncbi.nlm.nih.gov/pubmed/?term=17420468
Title:
URL: http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=gse2034; http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=gse3494
PMIDs: 15721472, 17420468
No abstract available.
```

```

notes:
  summary:
    15721472: A gene signature was identified that can accurately predict distant metastases in node-negative cases. 17420468: Tumors with a lung metastatic gene signature were shown to be larger.
  mapping.method:
    maxRowVariance
  mapping.group:
    EntrezGene.ID
  preprocessing:
    As published by original author.

featureData(eset):
An object of class 'AnnotatedDataFrame'
  featureNames: 1007_s_at 1053_at ... AFFX-HUMISGF3A/M97935_MB_at
    (21169 total)
  varLabels: probeset gene EntrezGene.ID best_probe
  varMetadata: labelDescription

```

Details

```

assayData: 21169 features, 344 samples
Platform type:
-----
Available sample meta-data:
-----

sample_name:
  Length      Class      Mode
    344 character character

alt_sample_name:
  Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
   3.0   122.8   605.5   575.7   836.5   2038.0

sample_type:
tumor
  344

er:
negative positive
   135       209

tumor_size:
  1    2    3    4 NA's
146 132   5   3   58

N:

```

```

0
344

age_at_initial_pathologic_diagnosis:
  Min. 1st Qu.  Median    Mean 3rd Qu.    Max.   NA's
  26.00   44.00   52.00   53.88   63.00   83.00    58

grade:
  1     2     3 NA's
  7    42   148  147

dmfs_days:
  Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
    61    1254    2616    2377    3285    5201

dmfs_status:
norecurrence  recurrence
          226           118

treatment:
untreated
      344

batch:
 VDX VDXN
 286   58

uncurated_author_metadata:
  Length    Class      Mode
    344 character character

```

Source

<http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=gse2034>; <http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=gse5327>

Index

* datasets

CAL, [3](#)
DFHCC, [5](#)
DFHCC2, [8](#)
DFHCC3, [10](#)
DUKE, [11](#)
DUKE2, [13](#)
duplicates, [15](#)
EMC2, [16](#)
EORTC10994, [18](#)
EXPO, [20](#)
FNCLCC, [22](#)
GSE25066, [23](#)
GSE32646, [26](#)
GSE48091, [28](#)
GSE58644, [29](#)
HLP, [32](#)
IRB, [34](#)
KOO, [36](#)
LUND, [40](#)
LUND2, [42](#)
MAINZ, [44](#)
MAQC2, [46](#)
MCCC, [48](#)
MDA4, [49](#)
METABRIC, [51](#)
MSK, [54](#)
MUG, [56](#)
NCCS, [57](#)
NCI, [59](#)
NKI, [61](#)
PNC, [64](#)
STK, [66](#)
STN02, [68](#)
TCGA, [71](#)
TRANSBIG, [73](#)
UCSF, [75](#)
UNC4, [78](#)
UNT, [81](#)

UPP, [83](#)
VDX, [85](#)

CAL, [3](#)

DFHCC, [5](#)
DFHCC2, [8](#)
DFHCC3, [10](#)
DUKE, [11](#)
DUKE2, [13](#)
duplicates, [15](#)

EMC2, [16](#)
EORTC10994, [18](#)
EXPO, [20](#)

FNCLCC, [22](#)

GSE25066, [23](#)
GSE32646, [26](#)
GSE48091, [28](#)
GSE58644, [29](#)

HLP, [32](#)

IRB, [34](#)

KOO, [36](#)

loadBreastDatasets, [38](#)

loadBreastEsets, [39](#)

LUND, [40](#)
LUND2, [42](#)

MAINZ, [44](#)
MAQC2, [46](#)
MCCC, [48](#)
MDA4, [49](#)
METABRIC, [51](#)
MSK, [54](#)
MUG, [56](#)

NCCS, [57](#)

NCI, [59](#)

NKI, [61](#)

PNC, [64](#)

STK, [66](#)

STN02, [68](#)

TCGA, [71](#)

TRANSBIG, [73](#)

UCSF, [75](#)

UNC4, [78](#)

UNT, [81](#)

UPP, [83](#)

VDX, [85](#)